

12-8082: Anti-Human IL-2R alpha (CD25) (Basiliximab) - PE

| | |
|--------------------------------|---|
| Clonality : | Monoclonal |
| Clone Name : | Hu107 |
| Application : | FACS |
| Alternative Name : | IL-2-RA; IL2-RA; TAC antigen; p55; CD25 |
| Isotype : | Human IgG1k |
| Immunogen Information : | Human CD25 |

Description

Expression Host : HEK-293

This non-therapeutic biosimilar antibody uses the same variable region sequence as the therapeutic antibody Basiliximab. Basiliximab recognizes human CD25. This product is for research use only.

CD25, a 55 kD type I transmembrane glycoprotein, has been shown to play roles in lymphocyte differentiation, activation, and proliferation. Many resting memory T cells constitutively express IL2Ralpha. It functions as the receptor for HTLV-1, resulting in its expression on neoplastic cells in adult T cell lymphoma/leukemia. CD25 (sIL-2R) has been used to track disease progression. Some additional clinical applications include Chagas disease, a disease characterized by a decline of CD25 expression on immune cells, and Multiple sclerosis, in which treatments with mAbs target CD25. Anti-Human IL-2R alpha (CD25) (Basiliximab) utilizes the same variable regions from the therapeutic antibody Basiliximab making it ideal for research projects.

Product Info

| | |
|----------------------------|---|
| Amount : | 50 µg |
| Content : | Concentration : 0.2 mg/ml This R-phycoerythrin (R-PE) conjugate is formulated in 0.01 M phosphate buffered saline (150 mM NaCl) PBS pH 7.4, 1% BSA and 0.09% sodium azide as a preservative. |
| Storage condition : | This R-phycoerythrin (R-PE) conjugate is stable when stored at 2-8°C. Do not freeze. |

Application Note

The suggested concentration for Basiliximab biosimilar antibody for staining cells in flow cytometry is ≤ 1.0 µg per 10⁶ cells in a volume of 100 µl. Titration of the reagent is recommended for optimal performance for each application.