

## 12-9147: Anti-SSTR2(tidutamab biosimilar) mAb

|                           |                                   |
|---------------------------|-----------------------------------|
| <b>Clonality :</b>        | Monoclonal                        |
| <b>Application :</b>      | ELISA                             |
| <b>Reactivity :</b>       | Human                             |
| <b>Alternative Name :</b> | SS-2-R, SS2-R, SS2R, SST2, SRIF-1 |
| <b>Isotype :</b>          | IgG1                              |

### Description

Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.

### Product Info

|                            |  |
|----------------------------|--|
| <b>Amount :</b>            | 50 µg / 100 µg   |
| <b>Purification :</b>      | Purified from cell culture supernatant by affinity chromatography  |
| <b>Content :</b>           | Not Sterile  |
| <b>Storage condition :</b> | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).Lyophilized proteins are shipped at ambient temperature. |

### Application Note

ELISA 1:5000-10000, FACS 1:100




Figure 1. ELISA plate pre-coated by 2 µg/mL (100 µL/well) Human SSTR2 Protein, hFc Tag can bind Anti-SSTR2 Neutralizing antibody in a linear range of 2.93-187.50 ng/mL. In order to specifically detect the Anti-SSTR2 Neutralizing antibody, mouse anti-human Fab-specific antibody was used as detection antibody.




Figure 2. Flow cytometry analysis with 1 µg/mL Anti-SSTR2 (tidutamab biosimilar) mAb on Expi293 cells transfected with Human SSTR2 protein (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).