

12-9230: Anti-NKG2D antibody(DMC287), IgG1 Chimeric mAb

Clonality :	Monoclonal
Clone Name :	DMC287
Application :	FACS
Reactivity :	Human
Alternative Name :	NKG2D,CD314,KLRK1,NK cell receptor D

Description

Natural killer (NK) cells are lymphocytes that can mediate lysis of certain tumor cells and virus-infected cells without previous activation. They can also regulate specific humoral and cell-mediated immunity. NK cells preferentially express several calcium-dependent (C-type) lectins, which have been implicated in the regulation of NK cell function. The NKG2 gene family is located within the NK complex, a region that contains several C-type lectin genes preferentially expressed in NK cells. This gene encodes a member of the NKG2 family. The encoded transmembrane protein is characterized by a type II membrane orientation (has an extracellular C terminus) and the presence of a C-type lectin domain. It binds to a diverse family of ligands that include MHC class I chain-related A and B proteins and UL-16 binding proteins, where ligand-receptor interactions can result in the activation of NK and T cells. The surface expression of these ligands is important for the recognition of stressed cells by the immune system, and thus this protein and its ligands are therapeutic targets for the treatment of immune diseases and cancers. Read-through transcription exists between this gene and the upstream KLRC4 (killer cell lectin-like receptor subfamily C, member 4) family member in the same cluster.

Product Info

Amount : Purification :	100 μg Purified from cell culture supernatant by affinity chromatography
Content :	Not Sterile
Storage condition :	Store at -20°C for 12 months (Avoid repeated freezing and thawing)

Application Note

FACS 1/100



Figure 1. Flow cytometry analysis with Anti-NKG2D (DMC287) on Expi293 cells transfected with human NKG2D (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).