

## 12-9258: Anti-B7-2 antibody(DM84), Rabbit mAb

Clonality :	Monoclonal
Clone Name :	DM84
Application :	ELISA,FACS
Reactivity :	Human
Alternative Name :	CD86, B7-2, B70, CD28LG2, LAB72, MGC34413,B72

## **Description**

This gene encodes a type I membrane protein that is a member of the immunoglobulin superfamily. This protein is expressed by antigen-presenting cells, and it is the ligand for two proteins at the cell surface of T cells, CD28 antigen and cytotoxic T-lymphocyte-associated protein 4. Binding of this protein with CD28 antigen is a costimulatory signal for activation of the T-cell. Binding of this protein with cytotoxic T-lymphocyte-associated protein 4 negatively regulates T-cell activation and diminishes the immune response. Alternative splicing results in several transcript variants encoding different isoforms.

## **Product Info**

100 µg
Purified from cell culture supernatant by affinity chromatography
Not Sterile
Store at -20°C for 12 months (Avoid repeated freezing and thawing)

**Application Note** 

ELISA 1/5000-10000;FACS 1/100




Figure 1. ELISA plate pre-coated by 2  $\hat{1}_{4}$ g/ml (100  $\hat{1}_{4}$ l/well) Human B7-2 protein, mFc-His tagged protein can bind Rabbit anti-B7-2 monoclonal antibody (clone: DM84) in a linear range of 1-100 ng/ml.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com




Figure 2. Expi 293 cell line transfected with irrelevant protein (A) and human B7-2 (B) were surface stained with Rabbit anti-B7-2 monoclonal antibody  $1\hat{1}$ /4g/ml (clone: DM84) followed by Alexa 488-conjugated anti-rabbit IgG secondary antibody.