

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

12-9266: Anti-BTN3A1 antibody(DM92), Rabbit mAb

Clonality: Monoclonal
Clone Name: DM92
Application: ELISA,FACS
Reactivity: Human

Alternative Name: BTN3A-1, BTF5, CD277, BTN3.1, BT3.1

Description

The butyrophilin (BTN) genes are a group of major histocompatibility complex (MHC)-associated genes that encode type I membrane proteins with 2 extracellular immunoglobulin (Ig) domains and an intracellular B30.2 (PRYSPRY) domain. Three subfamilies of human BTN genes are located in the MHC class I region: the single-copy BTN1A1 gene (MIM 601610) and the BTN2 (e.g., BTN2A1; MIM 613590) and BTN3 (e.g., BNT3A1) genes, which have undergone tandem duplication, resulting in 3 copies of each.

Product Info

Amount: 100 μg

Purification: Purified from cell culture supernatant by affinity chromatography

Content: Not Sterile

Storage condition: Store at -20°C for 12 months (Avoid repeated freezing and thawing)

Application Note

ELISA 1/5000-10000; FACS 1/100




Figure 1. ELISA plate pre-coated by 2 $1\frac{1}{4}$ g/ml (100 $1\frac{1}{4}$ l/well) Human BTN3A1 protein, mFc-His tagged protein can bind Rabbit anti-BTN3A1 monoclonal antibody (clone: DM92) in a linear range of 0.64-80 ng/ml.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2. Flow cytometry analysis with Anti-BTN3A1 (DM92) on Expi293 cells transfected with human BTN3A1 (Red histogram) or Expi293 transfected with irrelevant protein (Blue histogram).