

## 32-13664: sCD40L Human, His Active

**Format :** sCD40L protein (0.5mg/ml) contains 20mM Tris-HCl buffer (pH 8.0), 0.2M NaCl, 1mM DTT and 40% glycerol.

**Alternative Name :** CD40 ligand, CD154, CD40L, gp39, hCD40L, HIGM1, IGM, IMD3, T-BAM, TNFSF5, TRAP.

### Description

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered colorless solution.

Biological Activity: Measured by its binding ability in a functional ELISA with Human CD40/TNFRSF5.

CD40L or CD154 is a membrane glycoprotein and differentiation antigen expressed on the surface of T-cells. The CD40 ligand stimulates B-cell proliferation and secretion of all immunoglobulin isotypes in the presence of cytokines. CD40 ligand has been shown to induce cytokine production and tumoricidal activity in peripheral blood monocytes. It also costimulates proliferation of activated T-cells and this is accompanied by the production of IFN-gamma, TNF-alpha, and IL2.

sCD40L Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 169 amino acids (113-261a.a.) and having a molecular mass of 18.3kDa. sCD40L is fused to a 20 amino acid His tag at N-Terminus and purified by proprietary chromatographic techniques.

### Product Info

**Amount :** 10 µg / 2 µg

**Purification :** Greater than 95.0% as determined by SDS-PAGE.

**Storage condition :** Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

**Amino Acid :** MGSSHHHHH SSGLVPRGSH MQKGDQNPQI AAHVISEASS KTTSVLQWAE KGYTMSNNL  
VTLENGKQLT VKRQGLYYIY AQVTFCSNRE ASSQAPFIAS LCLKSPGRFE RILLRAANTH SAKPCGQQS  
IHLGGVFELQ PGASVFNVT DPSQVSHGTG FTSFGLLKL