

32-13724: C3a Human

Format : C3a solution contains Phosphate buffered saline, pH 7.2.

Alternative Name : Complement C3, C3 and PZP-like alpha-2-macroglobulin domain-containing protein 1, C3, CPAMD1.

Description

Source: Human Plasma.

Physical Appearance: Sterile Filtered solution.

Biological Activity: null

Native human C3a is created by cleavage of human C3 protein by a human C3 convertase. C3a is a member of the anaphylatoxin family of three proteins (C3a, C4a and C5a) created by the activation of complement. C3a mediates various inflammatory responses including smooth muscle contraction, increased vascular permeability, vasodilation and release of histamine from mast cells and basophils. C3a (In contrast to C5a) does not exhibit significant neutrophil stimulating activities and does not induce chemotaxis, granule release or superoxide production. C3a acts through the C3a Receptor which is a G-protein coupled receptor found mainly on peripheral tissues, lymphoid cells and in the central nervous system. These activities of C3a are inactivated by removal of the C-terminal arginine which happens rapidly in plasma due to the action of carboxypeptidase N.

Human Complement C3a produced in Human plasma is a non glycosylated polypeptide chain containing 77 amino acids having a molecular weight of 9,089 daltons.

Product Info

Amount : 50 µg / 10 µg

Purification : Greater than 98.0% as determined by SDS-PAGE.

Storage condition : C3a Human is stable at 4°C if entire vial will be used within 2-4 weeks. Store, frozen below -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.