

## 32-13737: C8 Human

**Format :** C8 protein solution contains PBS pH 7.2.

### Description

Source: Human Plasma.

Physical Appearance: Sterile filtered solution.

Biological Activity: null

Native human C8 is a glycosylated protein consist of 3 polypeptide chains: The alpha chain and the gamma chain are disulfide linked whereas, The beta chain is non-covalently bound to the  $\alpha$ -g complex. C8 is necessary for formation of the membrane attack complex and is activated by binding on the cell membrane newly-formed C5b, C6, C7 complexes. Each pathway of complement activation generates proteolytic enzyme complexes which bind the target surface. These enzymes cleave a peptide bond in the larger alpha chain of C5 releasing C5a and activating C5b. Although C5b is unstable it remains bound to the activating complex for a few minutes during which it binds a single C6 from the surrounding fluid or it decays and is no longer capable of forming MAC. The C5b,6 complex may also remain connected to the C3/C5 convertase where the binding of a single C7 exposes a membrane-binding region and C5b,6,7 can enter into the bilipid layer of the target cell. Each C5b-7 complex can bind 1 molecule of C8 causing the complex to enter more firmly into the membrane.

Human Complement C8 produced in Human plasma is glycosylated polypeptide chain containing 3 chains and having a total molecular mass of 151kDa.

### Product Info

**Amount :** 100  $\mu$ g / 20  $\mu$ g

**Purification :** Greater than 95% as determined by SDS-PAGE.

**Storage condition :** C8 Human is stable at 4°C if entire vial will be used within 2-4 weeks. Store, frozen below -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.