

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-13828: Borrelia p30

Format: Borrelia p30 is supplied in 20mM HEPES buffer pH-7.5, 400mM NaCl and 20% glycerol.

## **Description**

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered clear solution.

**Biological Activitynull** 

Borrelia is a part of the genus of bacteria of the spirochete phylum. Borrelia causes borreliosis, which is a zoonotic, vectorborne disease transmitted mostly by ticks and some by lice, depending on the species. Of the 36 known species of Borrelia, 12 are distinguished to cause Lyme disease or borreliosis and are transmitted by ticks. The Gram-negative bacterium Borreliella spielmanii is one of the pathogens of the Borreliella burgdorferi sensu lato complex causing Lyme disease. DbpA, also known as p17 or Osp17, is a heterogeneous protein among the human pathogenic B. burgdorferi sensu lato species. Borrelia Burgdorferi p30 Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain having a molecular mass of 29kDa. Borrelia p30 is expressed with a 6xHis tag and purified by proprietary chromatographic techniques.

## **Product Info**

Amount:  $20 \mu g / 5 \mu g$ 

**Purification:** Greater than 80.0% as determined by SDS-PAGE.

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods **Storage condition:** 

of time. Avoid multiple freeze-thaw cycles.