

30-2714: Anti-Hu TCR Vgamma4 PE

| | |
|--------------------------------|---------------|
| Clonality : | Monoclonal |
| Clone Name : | 4A11.904 |
| Application : | FACS |
| Reactivity : | Human |
| Conjugate : | PE |
| Alternative Name : | TCRVG4 |
| Immunogen Information : | human T cells |

Description

TCR Vgamma4 is a variant of TCR gamma chain, that is present on a minor subset of human gamma/delta T cells. Specificity :The mouse monoclonal antibody 4A11.904 recognizes an extracellular epitope of human TCR Vgamma4, that is present on a minor subset of human gamma/delta T cells.

Product Info

| | |
|----------------------------|---|
| Amount : | 100 Tests |
| Purification : | Purified antibody is conjugated with R-phycoerythrin (PE) under optimum conditions. Unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography. |
| Content : | Concentration: 1 mg/ml Storage Buffer: Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide |
| Storage condition : | Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze. |

Application Note

Flow cytometry: The reagent is designed for analysis of human blood cells using 10 μ l reagent / 100 μ l of whole blood or 10^6 cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.




Figure 1: Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human TCR Vgamma4 (4A11.904) PE antibody




Figure 2: Flow cytometry multicolor surface staining pattern of human lymphocytes stained using anti-human TCR Vgamma4 (4A11.904) PE antibody and anti-human CD3 (UCHT1) APC antibody




Figure 3: Separation of human TCR Vgamma4 positive lymphocytes (red-filled) from human TCR Vgamma4 negative lymphocytes (black-dashed) in flow cytometry analysis (surface staining) of peripheral whole blood stained using anti-human TCR Vgamma4 (4A11.904) PE antibody