

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 30-2731B: Biotin Conjugated Anti-Ms Ly6G (clone: RB6-8C5)

Clonality: Monoclonal
Clone Name: RB6-8C5
Application: IHC,FACS,WB
Reactivity: Mouse
Conjugate: Biotin
Gene: Ly6G
Gene ID: 546644

Uniprot ID: A0A087WQF5

Alternative Name: lymphocyte antigen 6 complex, locus G Gr1, Gr-1, Ly-6G

Immunogen Information: Murine granulocytes

## **Description**

Ly6G is a component of the myeloid differentiation antigen Gr-1, together with Ly6C. Ly6G is a good marker for detection of peripheral neutrophils. Expression of Gr-1 in bone marrow correlates with granulocyte differentiation and maturation. Physiological role of Ly6G remains still unclear. Its treatment with antibodies in vivo leads to neutropenia and has inhibitory effect on local immune responses.

Specificity: The rat monoclonal antibody RB6-8C5 detects Ly6G component of Gr-1 antigen, a commonly used surface marker of neutrophils.

## **Product Info**

Amount: 0.1 mg

Purified antibody is conjugated with biotin LC-NHS ester under optimum conditions and

unconjugated antibody and free biotin are removed by size-exclusion chromatography.

Content: Concentration: 1 mg/ml

Storage Buffer: Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide

**Storage condition :** Store at 2-8°C. Do not freeze.

## **Application Note**

Flow cytometry: Recommended dilution: 2-5 µg/ml.




Figure 1: Flow cytometry surface staining pattern of murine peritoneal fluid cell suspension stained using anti-murine Ly6G (RB6-8C5) Biotin antibody



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2: Separation of murine myeloid Ly6G positive cells stained using anti-Ly6G (RB-8C5) Biotin antibody (Streptavidin APC, red-filled) from murine myeloid cells unstained by primary antibody (Streptavidin APC, black-dashed) in flow cytometry analysis.