

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

30-2732: Anti-Hu LOX1 Purified

Clonality: Monoclonal **Clone Name:** 15C4 Application: ICC Human Reactivity: Gene: OLR1 Gene ID: 4973 **Uniprot ID:** P78380 Format: **Purified**

Alternative Name: oxidized low density lipoprotein receptor 1 OLR1, SLOX1, SCARE1, CLEC8A

Immunogen Information: fusion protein of human LOX1 ectodomain and of human IgG Fc

Description

LOX1, a 31 kDa type II transmembrane protein, is a C-type lectin, functioning as a scavenger receptor for e.g. oxidized low density lipoprotein, apoptotic heat shock proteins, or CRP. It is expressed by macrophages, fibroblasts, platelets, endothelial cells, and smooth muscle cells, and its defects can lead to atherosclerosis. Its expression is enhanced under inflammatory conditions. Multiple splicing variants have been identified.

Specificity: The mouse monoclonal antibody 15C4 recognizes an extracellular epitope of LOX1, a C-type lectin transmembrane protein.

Product Info

Amount: 0.1 mg

Purification : Purified by protein-A affinity chromatography.

Content: Concentration: 1 mg/ml

Storage Buffer: Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide

Storage condition : Store at 2-8°C. Do not freeze.

Application Note

Flow cytometry: Recommended dilution: 1-5 µg/ml




Figure 1: Flow cytometry surface staining pattern of human dendritic cells in flow cytometry analysis (surface staining) stained using anti-human LOX1 (15C4) purified antibody



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2: Separation of human dendritic cells (red-filled) from human lymphocytes (black-dashed) in flow cytometry analysis (surface staining) stained using anti-human LOX1 (15C4) purified antibody (GAM APC).