

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

30-2749: Anti-Ms CD8a FITC

Clonality: Monoclonal **Clone Name:** 53-6.7 Application: FACS Reactivity: Mouse Conjugate: FITC Gene: CD8a Gene ID: 12525 **Uniprot ID:** P01731

Alternative Name: CD8a molecule Ly-2, Ly-B, Ly-35, Lyt-2, BB154331

Immunogen Information: Murine spleen cells

Description

The CD8a (CD8 alpha) subunit of CD8 T cell coreceptor is expressed in CD8 alpha/beta heterodimers on majority of MHC I-restricted conventional T cells and thymocytes and in CD8 alpha/alpha homodimers on subsets of memory T cells, intraepithelial lymphocytes, NK cells, macrophages and dendritic cells. Regulation of CD8 beta level on T cell surface seems to be an important mechanism to control their effector function. Assembly of CD8 alpha/beta but not alpha/alpha dimers is connected with formation or localization to the lipid rafts. Recruiting triggered TCR complexes to these membrane microdomains as well as affinity of TCR to MHC I is modulated by CD8, thereby affecting the functional diversity of the TCR signaling.

Specificity: The rat monoclonal antibody 53-6.7 recognizes an extracellular epitope of murine CD8a (32-34 kDa; alpha chain of the CD8 antigen).

Product Info

Amount: 0.1 mg

Purified antibody is conjugated with fluorescein isothiocyanate (FITC) under optimum conditions

Purification: and unconjugated antibody and free fluorochrome are removed by size-exclusion

chromatography.

Content: Concentration: 0.5 mg/ml

Storage Buffer: Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide

Storage condition : Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.

Application Note

Flow cytometry: Recommended dilution: 1-2 µg/ml.