

30-2768: Anti-Hu CD52 APC

| | |
|--------------------------------|-------------------------------------|
| Clonality : | Monoclonal |
| Clone Name : | 4C8 |
| Application : | FACS |
| Reactivity : | Human |
| Conjugate : | APC |
| Gene : | CD52 |
| Gene ID : | 1043 |
| Uniprot ID : | P31358 |
| Alternative Name : | CD52 molecule CAMPATH-1, HE5, ESPE5 |
| Immunogen Information : | human lymphocytes |

Description

CD52 (CAMPATH-1, HE5) is a highly glycosylated GPI-anchored 21-28 kDa glycopeptide which is present at high levels on lymphocytes, macrophages, epithelial cells of male reproductive tract and mature sperm. Its 12-amino acid backbone carries a complex N-linked carbohydrate moiety, which differs between sperm and leukocyte CD52, as well as the GPI anchor does. CD52 can be acquired by sperm cells from seminal plasma, where it is released by epithelial cells. Although CD52 is not an essential T-cell costimulator, its triggering results in activation of normal human T cells. CD52 is a very good target for antibody/complement-mediated cell lysis.

Specificity : The mouse monoclonal antibody 4C8 recognizes CD52, a GPI-anchored glycopeptide expressed on the surface of lymphocytes, macrophages, epithelial cells of male reproductive tract, and mature sperms.

Product Info

| | |
|----------------------------|---|
| Amount : | 100 Tests |
| Purification : | Purified antibody is conjugated with activated allophycocyanin (APC) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography. |
| Content : | Storage Buffer: Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide |
| Storage condition : | Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze. |

Application Note

Flow cytometry: The reagent is designed for analysis of human blood cells using 10 μl reagent / 100 μl of whole blood or 10^6 cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.