

## 30-2780: Anti-Ms CD4 Purified

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	GK1.5
<b>Application :</b>	FACS, IP, IHC-Fr, ICC
<b>Reactivity :</b>	Mouse
<b>Gene :</b>	CD4
<b>Gene ID :</b>	12504
<b>Uniprot ID :</b>	P06332
<b>Format :</b>	Purified
<b>Alternative Name :</b>	CD4 molecule T4/Leu-3, L3T4
<b>Immunogen Information :</b>	Mouse CTL clone V4 cells

### Description

CD4 (T4) is a single chain transmembrane glycoprotein and belongs to immunoglobulin supergene family. In extracellular region there are 4 immunoglobulin-like domains (1 Ig-like V-type and 3 Ig-like C2-type). Transmembrane region forms 25 aa, cytoplasmic tail consists of 38 aa. Domains 1,2 and 4 are stabilized by disulfide bonds. The intracellular domain of CD4 is associated with p56Lck, a Src-like protein tyrosine kinase. It was described that CD4 segregates into specific detergent-resistant T-cell membrane microdomains. Extracellular ligands: MHC class II molecules (binds to CDR2-like region in CD4 domain 1); HIV envelope protein gp120 (binds to CDR2-like region in CD4 domain 1); IL-16 (binds to CD4 domain 3), human seminal plasma glycoprotein gp17 (binds to CD4 domain 1), L-selectin. Intracellular ligands: p56LckCD4 is a co-receptor involved in immune response (co-receptor activity in binding to MHC class II molecules) and HIV infection (human immunodeficiency virus; CD4 is primary receptor for HIV-1 surface glycoprotein gp120). CD4 regulates T-cell activation, T/B-cell adhesion, T-cell differentiation, T-cell selection and signal transduction. Defects in antigen presentation (MHC class II) cause dysfunction of CD4+ T-cells and their almost complete absence in patients blood, tissue and organs (SCID immunodeficiency).

Specificity :The rat monoclonal antibody GK1.5 reacts with an extracellular epitope of mouse CD4 transmembrane glycoprotein (55 kDa).

### Product Info

<b>Amount :</b>	0.1 mg
<b>Purification :</b>	Purified by protein-G affinity chromatography.
<b>Content :</b>	Concentration: 0.5 mg/ml Storage Buffer: Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
<b>Storage condition :</b>	Store at 2-8°C. Do not freeze.

### Application Note

Immunocytochemistry: Recommended dilution: 1-4  $\mu\text{g/ml}$ .

Immunoprecipitation: Recommended dilution: 1-2  $\mu\text{g}$  / 100-500  $\mu\text{g}$  of protein in 1 ml lysate.

Flow cytometry: Recommended dilution: 1  $\mu\text{g}$ /million cells.