

### 30-2785: Anti-Hu CD369 PE

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	15E2
<b>Application :</b>	FACS
<b>Reactivity :</b>	Human
<b>Conjugate :</b>	PE
<b>Gene :</b>	CLEC7A
<b>Gene ID :</b>	64581
<b>Uniprot ID :</b>	Q9BXN2
<b>Alternative Name :</b>	C-type lectin domain containing 7A Dectin-1, CLEC7A, BGR, CANDF4, SCARE2
<b>Immunogen Information :</b>	Extracellular part of human CD369 with hIgG FC tag

#### Description

CD369 (dectin-1, beta-glucan receptor) is a 33 kDa type II transmembrane glycoprotein of lectin family, and serves as a part of innate immunity system by binding to beta-glucan polymers, which are typical for yeast and mycobacterial cell walls. CD369 is expressed predominantly on dendritic cells, but it can be detected also on monocytes, macrophages, mast cells, eosinophils, B cells, endothelial cells, and sometimes also on some T cell subsets.

Specificity :The mouse monoclonal antibody 15E2 recognizes an extracellular epitope LWEDGSTFSSN of human CD369 (Dectin-1), a 33 kDa transmembrane glycoprotein expressed predominantly on dendritic cells.

#### Product Info

<b>Amount :</b>	100 Tests
<b>Purification :</b>	Purified antibody is conjugated with R-phycoerythrin (PE) under optimum conditions. Unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.
<b>Content :</b>	Concentration: 1 mg/ml Storage Buffer: Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
<b>Storage condition :</b>	Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.

#### Application Note

Flow cytometry: The reagent is designed for analysis of human blood cells using 10  $\mu$ l reagent / 100  $\mu$ l of whole blood or  $10^6$  cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.