

## 30-2806: Anti-Hu CD3 Biotin

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	UCHT1
<b>Application :</b>	FACS, IP, IHC-Fr
<b>Reactivity :</b>	Non-human primates, Human
<b>Conjugate :</b>	Biotin
<b>Gene :</b>	CD3E
<b>Gene ID :</b>	916
<b>Uniprot ID :</b>	P07766
<b>Alternative Name :</b>	CD3 antigen, epsilon polypeptide CD3E, T3E, TCRE
<b>Immunogen Information :</b>	human thymocytes followed by Sezary T cells

### Description

CD3 complex is crucial in transducing antigen-recognition signals into the cytoplasm of T cells and in regulating the cell surface expression of the TCR complex. T cell activation through the antigen receptor (TCR) involves the cytoplasmic tails of the CD3 subunits CD3 gamma, CD3 delta, CD3 epsilon and CD3 zeta. These CD3 subunits are structurally related members of the immunoglobulins super family encoded by closely linked genes on human chromosome 11. The CD3 components have long cytoplasmic tails that associate with cytoplasmic signal transduction molecules. This association is mediated at least in part by a double tyrosine-based motif present in a single copy in the CD3 subunits. CD3 may play a role in TCR-induced growth arrest, cell survival and proliferation. The CD3 antigen is present on 68-82% of normal peripheral blood lymphocytes, 65-85% of thymocytes and Purkynje cells in the cerebellum. It is never expressed on B or NK cells. Decreased percentages of T lymphocytes may be observed in some autoimmune diseases.

**Specificity :** The antibody UCHT1 recognizes an extracellular epitope on CD3 antigen of the TCR/CD3 complex on mature human T cells. The UCHT1 antibody reacts with the epsilon chain of the CD3 complex.

### Product Info

<b>Amount :</b>	0.1 mg
<b>Purification :</b>	Purified antibody is conjugated with biotin LC-NHS ester under optimum conditions and unconjugated antibody and free biotin are removed by size-exclusion chromatography.
<b>Content :</b>	Storage Buffer: Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
<b>Storage condition :</b>	Store at 2-8°C. Do not freeze.

### Application Note

Flow cytometry: Extracellular and intracellular staining; recommended dilution: 1-12  $\mu\text{g/ml}$ .