

### 30-2809: Anti-Hu CD3 PE

|                                |  |
|--------------------------------|--|
| <b>Clonality :</b>             | Monoclonal                                       |
| <b>Clone Name :</b>            | UCHT1  |
| <b>Application :</b>           | FACS   |
| <b>Reactivity :</b>            | Non-human primates, Human                        |
| <b>Conjugate :</b>             | PE   |
| <b>Gene :</b>                  | CD3E   |
| <b>Gene ID :</b>               | 916  |
| <b>Uniprot ID :</b>            | P07766   |
| <b>Alternative Name :</b>      | CD3 antigen, epsilon polypeptide CD3E, T3E, TCRE |
| <b>Immunogen Information :</b> | human thymocytes followed by Sezary T cells      |

#### Description

CD3 complex is crucial in transducing antigen-recognition signals into the cytoplasm of T cells and in regulating the cell surface expression of the TCR complex. T cell activation through the antigen receptor (TCR) involves the cytoplasmic tails of the CD3 subunits CD3 gamma, CD3 delta, CD3 epsilon and CD3 zeta. These CD3 subunits are structurally related members of the immunoglobulins super family encoded by closely linked genes on human chromosome 11. The CD3 components have long cytoplasmic tails that associate with cytoplasmic signal transduction molecules. This association is mediated at least in part by a double tyrosine-based motif present in a single copy in the CD3 subunits. CD3 may play a role in TCR-induced growth arrest, cell survival and proliferation. The CD3 antigen is present on 68-82% of normal peripheral blood lymphocytes, 65-85% of thymocytes and Purkynje cells in the cerebellum. It is never expressed on B or NK cells. Decreased percentages of T lymphocytes may be observed in some autoimmune diseases.

**Specificity :** The antibody UCHT1 recognizes an extracellular epitope on CD3 antigen of the TCR/CD3 complex on mature human T cells. The UCHT1 antibody reacts with the epsilon chain of the CD3 complex.

#### Product Info

|                            |   |
|----------------------------|---|
| <b>Amount :</b>            | 100 Tests   |
| <b>Purification :</b>      | Purified antibody is conjugated with R-phycoerythrin (PE) under optimum conditions. Unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography. |
| <b>Content :</b>           | Concentration: 1 mg/ml<br>Storage Buffer: Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide   |
| <b>Storage condition :</b> | Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.  |

#### Application Note

**Flow cytometry:** The reagent is designed for analysis of human blood cells using 20  $\mu\text{l}$  reagent / 100  $\mu\text{l}$  of whole blood or  $10^6$  cells in a suspension. The content of a vial (2 ml) is sufficient for 100 tests. Extracellular and intracellular staining.