

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

30-2823: Anti-Hu CD172g PE

Clonality: Monoclonal Clone Name: OX-119 **FACS** Application: Reactivity: Human Conjugate: PΕ Gene: SIRPG Gene ID: 55423 **Uniprot ID:** 09P1W8

Alternative Name: signal regulatory protein gamma SIRP gamma, SIRP-B2

Immunogen Information: recombinant human CD172g

Description

CD172g is a transmembrane glycoprotein, which may play a role in inter-T cellular signaling by binding CD47, and thus in influencing T cell behaviour. CD172g is expressed on mature thymocytes, CD4+ T cells, CD8+ T cells, NK cells, and some B cells. It is absent on myeloid cells. Engagement of CD172g by CD47 expressed on antigen presenting cells results in enhanced antigen-specific T cell proliferation and costimulates T cell activation.

Specificity: The mouse monoclonal antibody OX-119 recognizes an extracellular epitope on CD172g, an approximately 55 kDa transmembrane glycoprotein expressed on most T cells, as well as on NK cells and some B cell populations.

Product Info

Amount: 100 Tests

Purification : Purified antibody is conjugated with R-phycoerythrin (PE) under optimum conditions.

Unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.

Concentration: 1 mg/ml

Content: Storage Buffer: Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide

Storage condition: Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.

Application Note

Flow cytometry: The reagent is designed for analysis of human blood cells using 10 $\tilde{A} \square \hat{A} \mu l$ reagent / 100 $\tilde{A} \square \hat{A} \mu l$ of whole blood or 10⁶ cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.