

30-2841: Anti-Human CD307b PE MAb (Clone: B24)

| | |
|--------------------------------|--|
| Clonality : | Monoclonal |
| Clone Name : | B24 |
| Application : | FACS |
| Reactivity : | Human |
| Conjugate : | PE |
| Gene : | FCRL2 |
| Gene ID : | 79368 |
| Uniprot ID : | Q96LA5 |
| Alternative Name : | FcRL2, FcRH2, IFGP4, IRTA4, SPAP1, Fc receptor like 2 |
| Isotype : | Mouse IgG2a kappa |
| Immunogen Information : | DNA-immunization followed by a boost with CD307b transfected cells |

Description

Specificity: The mouse monoclonal antibody B24 recognizes an extracellular epitope of CD307b, a transmembrane glycoprotein expressed mainly in mature and memory B cells.

CD307b is a type I transmembrane glycoprotein of the Fc receptor family. It contains one ITAM motif and two ITIM motifs in its cytoplasmic domain. It is expressed in spleen and lymph nodes in mature B cells and memory B cells. CD307b may be a prognostic marker for chronic lymphocytic leukemia.

Product Info

| | |
|----------------------------|---|
| Amount : | 100 Tests |
| Purification : | Purified antibody is conjugated with R-phycoerythrin (PE) under optimum conditions. Unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography. |
| Content : | Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide |
| Storage condition : | Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze. |

Application Note

Flow cytometry: The reagent is designed for analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10⁶ cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.




Fig 1 : Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human CD307b (B24) PE antibody (10 µl reagent / 100 µl of peripheral whole blood).




Fig 2 : Flow cytometry multicolor surface staining of human lymphocytes stained using anti-human CD307b(B24) PE antibody (10 μ l reagent / 100 μ l of peripheral whole blood) and anti-human CD19 (LT19) APC antibody (10 μ l reagent / 100 μ l of peripheral whole blood).




Fig 3 : Separation of human CD19 positive B cells (red-filled) from neutrophil granulocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human CD307b (B24) PE antibody (10 μ l reagent / 100 μ l of peripheral whole blood).