

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

30-2867: Anti-Human CD158z PE MAb(Clone :CH21)

Clonality: Monoclonal

Clone Name: CH21
Application: FACS
Reactivity: Human
Conjugate: PE
Gene: KIR3DL3
Gene ID: 115653
Uniprot ID: Q8N743

Alternative Name: killer cell immunoglobulin like receptor, three Ig, KIR3DL3

Isotype: Mouse IgG2a

Immunogen Information: human CD158z transfectants

Description

Specificity: The mouse monoclonal antibody CH21 recognizes an extracellular epitope of human CD158z (KIR3DL3), a transmembrane glycoprotein of killer cell inhibitory receptor family.

CD158z (KIR3DL3)is one of killer cell inhibitory receptors. It has three extracellular immunoglobulin-like domains and a long cytoplasmic tail, which, however, contains only one ITIM. Like other KIRs, CD158z is highly polymorphic, but it seems that its immunoglobulin-like domains are quite conserved among high primates.

Product Info

Amount: 0.1 mg

Purification: Purified antibody is conjugated with R-phycoerythrin (PE) under optimum conditions.

Unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.

Content: 0.1 mg/ml

Formulation: Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide

Storage condition : Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze. Stabilizing phosphate

buffered saline (PBS), pH 7.4, 15 mM sodium azide

Application Note

Flow cytometry: Recommended dilution: 1-5 µg/ml



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com




Fig 1: Flow cytometry surface staining patterns of KIR3DL3 (CD158z) transfected HEK-293 suspension stained using anti-human CD158z (CH21) PE antibody (concentration in sample 5 $\hat{l}\frac{1}{4}$ g/ml, left) or mouse IgG2a isotype control (MOPC-173) PE antibody (concentration in sample 5 $\hat{l}\frac{1}{4}$ g/ml, same as CD158z PE concentration, right).




Fig 2: Separation of CD158z positive cells (red-filled) from CD158z negative cells (black-dashed) in flow cytometry analysis (surface staining) of KIR3DL3 (CD158z) transfected HEK-293 suspension stained using anti-human CD158z (CH21) PE antibody (concentration in sample 5 $\hat{1}\frac{1}{4}$ g/ml).