

## 30-2890: Anti-Hu CD172ab APC Mab(SE5A5)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	SE5A5
<b>Application :</b>	FACS
<b>Reactivity :</b>	Human
<b>Conjugate :</b>	APC
<b>Alternative Name :</b>	PTPNS1, BIT, MFR, SIRPA, SHPS1, SIRPB1,
<b>Isotype :</b>	Mouse IgG1
<b>Immunogen Information :</b>	NIH-3T3 / human CD172a cell line

### Description

Specificity: The mouse monoclonal antibody SE5A5 recognizes a common extracellular epitope on human CD172a and CD172b antigens (approx. 90 kDa and approx. 50 kDa, respectively), although its reactivity with CD172a is higher.

CD172a, the signal-regulatory protein alpha (SIRP alpha), also known as SH2 domain-containing phosphatase substrate-1 (SHPS1), is a 75-110 kDa transmembrane glycoprotein expressed mainly on granulocytes, monocytes, macrophages, dendritic cells and neurons. Its extracellular ligand is CD47. CD172a serves as a substrate of activated receptor tyrosine kinases and upon phosphorylation it recruits SH2 domain-containing tyrosine phosphatases, thereby regulating signal transduction processes related to cell activation, transmigration and phagocytosis. CD172a is a specific marker of cardiomyocytes derived from human pluripotent stem cells and serves as a negative regulator of signaling and growth in myeloid progenitor cells. Extracellular part of CD172b is 90% identical to that of CD172a, but unlike CD172, it has dramatically reduced intracellular domain.

### Product Info

<b>Amount :</b>	100 Tests
<b>Purification :</b>	Purified antibody is conjugated with activated allophycocyanin (APC) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.
<b>Content :</b>	Formulation: Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
<b>Storage condition :</b>	Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.

### Application Note

Flow cytometry: The reagent is designed for analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10<sup>6</sup> cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.




Figure 1: Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human CD172ab (SE5A5) APC antibody (10  $\mu$ l reagent / 100  $\mu$ l of peripheral whole blood).




Figure 2: Separation of human monocytes (red-filled) from lymphocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human CD172ab (SE5A5) APC antibody (10  $\mu$ l reagent / 100  $\mu$ l of peripheral whole blood).