

30-2892: Anti-Hu CD68 FITC Mab(Y1/82A)

Clonality :	Monoclonal
Clone Name :	Y1/82A
Application :	FACS
Reactivity :	Human
Conjugate :	FITC
Gene ID :	968
Uniprot ID :	P34810
Alternative Name :	GP110, LAMP4, SCARD1
Isotype :	Mouse IgG2b
Immunogen Information :	Lysosomal contents of lung macrophages

Description

Specificity:The mouse monoclonal antibody Y1/82A recognizes CD68 (LAMP4), a 110 kDa glycoprotein expressed mainly in cytoplasmic granules of monocytes/macrophages, granulocytes, and dendritic cells.

CD68 (also known as LAMP4 or SCARD1) is a 110 kDa type I transmembrane glycoprotein of the lysosomal/endosomal-associated membrane glycoprotein (LAMP) family and the scavenger receptor family. Although CD68 primarily localizes to lysosomes and endosomes, its fraction circulates to the cell surface. By the heavily glycosylated extracellular domain CD68 binds to tissue- and organ-specific lectins or selectins. It is expressed mainly in cytoplasmic granules of monocytes/macrophages, granulocytes, and dendritic cells, but also e.g. in a proportion of epithelial tumours (diagnosis of poorly differentiated neoplasms).

Product Info

Amount :	100 Tests
Purification :	Purified antibody is conjugated with fluorescein isothiocyanate (FITC) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.
Content :	Formulation: Stabilizing phosphate-buffered saline (PBS), pH 7.4, 15 mM sodium azide
Storage condition :	Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.

Application Note

Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10⁶ cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests. Extracellular and intracellular staining.




Figure 1: Flow cytometry intracellular staining pattern of human peripheral whole blood stained using anti-human CD68 (Y1/82A) PE antibody (4 µl reagent / 100 µl of peripheral whole blood).




Figure 2: Separation of human monocytes (red-filled) from lymphocytes (black-dashed) in flow cytometry analysis (intracellular staining) of human peripheral whole blood stained using anti-human CD68 (Y1/82A) PE antibody (4 µl reagent / 100 µl of peripheral whole blood).