∗ abeomics

30-2897: Anti-Hu CD316 PE Mab(8A12)

Clonality :	Monoclonal
Clone Name :	8A12
Application :	FACS
Reactivity :	Human
Conjugate :	PE
Gene ID :	93185
Uniprot ID :	Q969P0
Alternative Name :	EWI2, PGRL, KCT-4, LIR-D1, IGSF8
Isotype :	Mouse IgG2a
Immunogen Information : human leukocytes	

Description

Specificity: The mouse monoclonal antibody 8A12 recognizes an extracellular epitope of CD316, a type I transmembrane glycoprotein, expressed mainly by lymphocytes, NK cells and in a broad range of tissues.

CD316 is an approximately 63 kDa transmembrane glycoprotein, which forms stable complexes with CD9, as well as with CD81. It seems that CD316 participates in regulation of cell polarity and motility, in cell proliferation, oocyte fertilization and tumor cell metastasis. It is expressed mainly by lymphocytes, NK cells and in a broad range of tissues, and associates with actin cytoskeleton.

Product Info

100 Tests
Purified antibody is conjugated with R-phycoerythrin (PE) under optimum conditions. Unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.
Formulation : Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.

Application Note

Flow cytometry: The reagent is designed for analysis of human blood cells using 10 μ l reagent / 100 μ l of whole blood or 10⁶ cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.



Figure 1: Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human CD316 (8A12) PE antibody (10 $\hat{1}$ /4l reagent / 100 $\hat{1}$ /4l of peripheral whole blood).



₩ abeomics

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com



Figure 2: Separation of human monocytes (red-filled) from neutrophil granulocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human CD316 (8A12) PE antibody (10 $\hat{1}$ /4l reagent / 100 $\hat{1}$ /4l of peripheral whole blood).