

## 36-3820-BSA-Azide-Free: Anti-CD134, Mouse (OX40) Monoclonal Antibody(Clone: OX-86) BSA/Azide Free

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	OX-86
<b>Application :</b>	FACS,IF
<b>Reactivity :</b>	Mouse
<b>Gene ID :</b>	22163
<b>Uniprot ID :</b>	P47741
<b>Alternative Name :</b>	CD134; IMD16; Lymphoid activation antigen (ACT35); OX40 cell surface antigen; OX40L receptor; TAX transcriptionally activated glycoprotein 1 receptor; Tumor necrosis factor receptor superfamily member 4 (TNFRSF4); Txgp1l
<b>Isotype :</b>	Rat IgG1, kappa
<b>Immunogen Information :</b>	Purified recombinant Mouse OX-40 antigen

### Description

CD134 is a type I integral membrane protein. This receptor is expressed on activated CD4+and CD8+T cells and B cells. The CD134 binds to CD134 ligand (CD252) to provide a costimulatory signal that is independent of CD28. CD134 is involved in coordinating CD4 T cell selection, migration and cytokine differentiation in T helper (Th)1 and Th2 cells. CD134 is also involved in the stimulation of T cells, T dependent humoral response and generation of optimal CD4+ T cell responses in vivo and in vitro. CD134 is expressed on activated CD4+ T lymphocytes, and its ligand, CD134L, is found preferentially on activated B cells. Engagement of CD134 with its ligand, CD134L, delivers a strong costimulatory signal to effector T cells.

### Product Info

<b>Amount :</b>	100 µg
<b>Content :</b>	1.0 mg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

### Application Note

Does not block binding of OX40L; Flow Cytometry (0.5-1ug/million cells in 0.1ml); Immunofluorescence (1-2ug/ml);




Fig. 1: SDS-PAGE Analysis Purified CD134 Rat Monoclonal Antibody (OX-86). Confirmation of Integrity and Purity of Antibody.