

30-2916: Anti-Hu CD75 PE Mab (LN1)

Clonality :	Monoclonal
Clone Name :	LN1
Application :	FACS
Reactivity :	Human
Conjugate :	PE
Alternative Name :	lactosamine
Isotype :	Mouse IgM kappa
Immunogen Information :	Stimulated human PBL

Description

Specificity: The mouse monoclonal antibody LN1 recognizes CD75, a lactosamine structure present mainly on the surface of B cell types.

CD75 is a lactosamine structure, which is present mainly on the surface of germinal center B cells. With lower level it is present on other mature B cells, and it is downregulated during differentiation to plasma cells. It is a marker which differentiates between malignant B cell types.

Product Info

Amount :	100 Tests
Purification :	Purified antibody is conjugated with R-phycoerythrin (PE) under optimum conditions. Unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.
Content :	Storage Buffer: Stabilizing Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide
Storage condition :	Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.

Application Note

Flow cytometry: The reagent is designed for analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10⁶ cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.




Figure 1: Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human CD75 (LN1) PE antibody (10 µl reagent / 100 µl of peripheral whole blood).




Figure 2: Separation of human CD75 positive lymphocytes (red-filled) from CD75 negative lymphocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human CD75 (LN1) PE antibody (10 μ g/ml reagent / 100 μ l of peripheral whole blood).