

## 32-18438: Human CCL17 Protein, hFc Tag

**Uniprot ID :** Q92583

**Alternative Name :** TARC; ABCD-2; SCYA17; A-152E5.3

### Description

**Description :** Recombinant human CCL17 Protein with C-terminal human Fc tag

**Background :** This antimicrobial gene is one of several Cys-Cys (CC) cytokine genes clustered on the q arm of chromosome 16. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for T lymphocytes, but not monocytes or granulocytes. The product of this gene binds to chemokine receptors CCR4 and CCR8. This chemokine plays important roles in T cell development in thymus as well as in trafficking and activation of mature T cells.

**Molecular Characterization:** mass of 34.2 kDa after removal of the signal peptide. The apparent molecular mass of CCL17-hFc is approximately 35-55 kDa due to glycosylation.

**Tag :** C-Human Fc tag

### Product Info

**Amount :** 50 µg / 100 µg

**Purification :** The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.

**Content :** Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization.

**Storage condition :** Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.




Figure 1. Human CCL17 Protein, hFc Tag on SDS-PAGE under reducing condition.