

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

11-3025: Polyclonal Antibody to mTLR3

Clonality: Polyclonal Application: FACS,WB Reactivity: Mouse Gene: Tlr3 Gene ID: 142980 **Uniprot ID:** 099MB1 Format: Purified **Alternative Name:** Tlr3

Isotype: Rabbit IgG

Immunogen Information: A partial length recombinant mTLR3 protein (amino acids 180-400) was used as the

immunogen for this antibody.

Description

Mouse Toll-like receptor 3 (mTLR3) belongs to a family of evolutionary conserved innate immune recognition molecules and recognizes double-stranded RNA, a molecular pattern associated with viral infections. It has seven exons. mTlr3 show high homology to hTLR3 in their predicted cDNA and protein sequences. The intracellular region of mTlr3 has remarkable homology with the same region of others mTlrs. mTlr3 mRNA is ubiquitously expressed in tissues, being expressed highest in spleen, kidney and lung. Its activation by the synthetic ligand polyinosine:polycytidylic acid (poly I:C) or by mRNA rapidly causes growth cone collapse and irreversibly inhibits neurite extension independent of NF-kappaB. The high levels of mTlr3 expression in Mz B cells perhaps contribute to their ability to clear antigen from the blood.

Product Info

Amount : 25 μg / 100 μg

Purification: Protein A Chromatography

Content: 25 µg in 50 µl/100 µg in 200 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium

azide is highly toxic.

Storage condition : Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

Western blot analysis: 2-4 μg/ml, Flowcytometric analysis: 1-2 μg/10⁶ Cells




Fig-1: Western blot analysis of mTLR3. Anti- mTLR3 antibody (11-3025) was used at 2 μ g/ml on EL-4 lysate.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Fig-2: Intracellular flowcytometric staining of mTLR3 in Splenocytes using 1 μ g/10^6 cells of Anti-mTLR3 (11-3025). Green represent Isotype control and red represent Anti-mTLR3 antibody (Abeomics). Goat Anti-Rabbit IgG FITC conjugated was used as the secondary antibody.