abeomics

9853 Pacific Heights Blvd.
Suite D. San Diego,
CA 92121, USA
Tel: 858-263-4982
Email: info@abeomics.com

12-9426: Anti-Nectin4 antibody(DMC438); IgG1 Chimeric mAb

Clonality :	Monoclonal
Clone Name :	DMC438
Application :	FACS
Reactivity :	Human
Uniprot ID :	Q96NY8
Alternative Name :	EDSS1; LNIR; nectin-4; PRR4; PVRL4
Isotype :	Rabbit/Human Fc chimeric IgG1

Description

Description :Anti-Nectin4 antibody(DMC438); IgG1 Chimeric mAb
This gene encodes a member of the nectin family. The encoded protein contains two immunoglobulin-like (Ig-like) C2-type domains and one Ig-like V-type domain. It is involved in cell adhesion through trans-homophilic and -heterophilic interactions. It is a single-pass type I membrane protein. The soluble form is produced by proteolytic cleavage at the cell surface by the metalloproteinase ADAM17:TACE. The secreted form is found in both breast tumor cell lines and breast tumor patients. Mutations in this gene are the cause of ectodermal dysplasia-syndactyly syndrome type 1; an autosomal recessive disorder. Alternatively spliced transcript variants have been found but the full-length nature of the variant has not been determined.

Product Info

|	
Amount :	10 µg / 100 µg
Purification :	Purified from cell culture supernatant by affinity chromatography
Content :	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage condition :	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

Application Note

FACS 1:100

Figure 1. Flow cytometry analysis with Anti-Nectin4 (DMC438) on Expi293 cells transfected with human Nectin4 (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).