

## 12-9436: Anti-FGFR4 antibody(DMC460); IgG1 Chimeric mAb

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	DMC460
<b>Application :</b>	FACS
<b>Reactivity :</b>	Human
<b>Uniprot ID :</b>	P22455
<b>Alternative Name :</b>	CD334; JTK2; TKF
<b>Isotype :</b>	Rabbit/Human Fc chimeric IgG1

### Description

Description :Anti-FGFR4 antibody(DMC460); IgG1 Chimeric mAb

The protein encoded by this gene is a tyrosine kinase and cell surface receptor for fibroblast growth factors. The encoded protein is involved in the regulation of several pathways; including cell proliferation; cell differentiation; cell migration; lipid metabolism; bile acid biosynthesis; vitamin D metabolism; glucose uptake; and phosphate homeostasis. This protein consists of an extracellular region; composed of three immunoglobulin-like domains; a single hydrophobic membrane-spanning segment; and a cytoplasmic tyrosine kinase domain. The extracellular portion interacts with fibroblast growth factors; setting in motion a cascade of downstream signals; ultimately influencing mitogenesis and differentiation. [provided by RefSeq; Aug 2017]

### Product Info

<b>Amount :</b>	10 µg / 100 µg
<b>Purification :</b>	Purified from cell culture supernatant by affinity chromatography
<b>Content :</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage condition :</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

### Application Note

FACS 1:100




Figure 1. Flow cytometry analysis with Anti-FGFR4 (DMC460) on Expi293 cells transfected with human FGFR4 (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).