

12-9439: Anti-CD142 antibody(DMC463); IgG1 Chimeric mAb

| | |
|---------------------------|--------------------------------|
| Clonality : | Monoclonal |
| Clone Name : | DMC463 |
| Application : | FACS |
| Reactivity : | Human |
| Uniprot ID : | P13726 |
| Alternative Name : | TF; Coagulation factor III; F3 |
| Isotype : | Rabbit/Human Fc chimeric IgG1 |

Description

Description :Anti-CD142 antibody(DMC463); IgG1 Chimeric mAb

This gene encodes coagulation factor III which is a cell surface glycoprotein. This factor enables cells to initiate the blood coagulation cascades; and it functions as the high-affinity receptor for the coagulation factor VII. The resulting complex provides a catalytic event that is responsible for initiation of the coagulation protease cascades by specific limited proteolysis. Unlike the other cofactors of these protease cascades; which circulate as nonfunctional precursors; this factor is a potent initiator that is fully functional when expressed on cell surfaces; for example; on monocytes. There are 3 distinct domains of this factor: extracellular; transmembrane; and cytoplasmic. Platelets and monocytes have been shown to express this coagulation factor under procoagulatory and proinflammatory stimuli; and a major role in HIV-associated coagulopathy has been described. Platelet-dependent monocyte expression of coagulation factor III has been described to be associated with Coronavirus Disease 2019 (COVID-19) severity and mortality. This protein is the only one in the coagulation pathway for which a congenital deficiency has not been described. Alternate splicing results in multiple transcript variants.[provided by RefSeq; Aug 2020]

Product Info

| | |
|----------------------------|--|
| Amount : | 10 µg / 100 µg |
| Purification : | Purified from cell culture supernatant by affinity chromatography |
| Content : | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Storage condition : | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |

Application Note

FACS 1:100




Figure 1. Flow cytometry analysis with Anti-CD142 (DMC463) on Expi293 cells transfected with human CD142 (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).