

## 36-1777: Monoclonal Antibody to Tyrosinase (Melanoma Marker)(Clone : SPM360)

Clonality :	Monoclonal
Clone Name :	SPM360
Application :	FACS,IF,IHC
Reactivity :	Human
Gene :	TYR
Gene ID :	7299
Uniprot ID :	P14679
Format :	Purified
Alternative Name :	TYR
Isotype :	Mouse IgG2a, kappa
Immunogen Information :	Recombinant tyrosinase protein

## Description


Recognizes a cluster of proteins between 70-80kDa, identified as tyrosinase. Occasionally a minor band at 55kDa is also detected. This MAb shows no cross-reaction with MAGE-1 and tyrosinase-related protein 1, TRP-1/gp75. Tyrosinase is a copper-containing metalloglycoprotein that catalyzes several steps in the melanin pigment biosynthetic pathway; the hydroxylation of tyrosine to L-3,4-dihydroxy-phenylalanine (dopa), and the subsequent oxidation of dopa to dopaquinone. Mutations of the tyrosinase gene occur in various forms of albinism. Tyrosinase is one of the targets for cytotoxic T-cell recognition in melanoma patients. Staining of melanomas with this MAb shows tyrosinase in melanotic as well as amelanotic variants. This MAb is a useful marker for melanocytes and melanomas.

## **Product Info**

Amount : Purification :	100 μg Affinity Chromatography
Content :	100 $\mu g$ in 500 $\mu l$ PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic.
Storage condition :	Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid repeated freeze and thaw cycles.

## **Application Note**

Flow Cytometry (0.5-1ug/million cells); Immunofluorescence (1-2ug/ml); ,Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 & degC followed by cooling at RT for 20 minutes),



Formalin-fixed, paraffin-embedded human Melanoma stained with Tyrosinase Monoclonal Antibody (SPM360).

For Research Use Only. Not for use in diagnostic/therapeutics procedures.