

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

30-1112PC-Cy5.5: Anti-CD7 Monoclonal Antibody (Clone:MEM-186) PerCP-Cy™5.5

Clonality: Monoclonal **Clone Name:** MEM-186 FACS Application: Reactivity: Human Gene: CD7 Gene ID: 924 **Uniprot ID:** P09564 Format: **Purified Alternative Name:** CD7

Isotype: Mouse IgG1

Immunogen Information: Human acute myelogenous leukaemia cell line KG-1.

Description

CD7, also known as gp40, is a member of the immunoglobulin superfamily found on T cells, NK cells, thymocytes, hematopoietic progenitors, and monocytes (weakly). CD7 is also expressed on acute lymphocytic leukemia (ALL). CD7 crosslinking induces a calcium flux in T lymphocytes, presumably as a result of cytoplasmic domain association with Pl3-kinase. CD7 co-stimulation can induce cytokine secretion and modulate cellular adhesion. A ligand of CD7, epithelial cell secreted protein K12, is produced in thymus to regulate thymocyte signaling and cytokine release. In lung microvascular endothelial cells CD7 serves as an IgM Fc receptor. Expression of CD7 is an important marker used in leukemia diagnostics.

Product Info

Amount: 100 tests

Purification: Purified by protein-A affinity chromatography

Storage condition : Store at 2-8°C. Do not freeze.

Application Note

Flow Cytometry Recommended dilution: 2 µg/ml

Immunoprecipitation Western Blotting Application note: Non-reducing conditions. Immunohistochemistry

Recommended dilution: 5 µg/ml

Positive tissue: spleen




Figure 1: Flow cytometry analysis (surface staining) of human peripheral blood cells with anti-human CD7 (MEM-186) PerCP-Cy™5.5.