

30-1242PE-Cy5: Anti-CD38 Monoclonal Antibody (Clone:HIT2) PE-Cy™ 5

Clonality :	Monoclonal
Clone Name :	HIT2
Application :	FACS
Reactivity :	Human
Gene :	CD38
Gene ID :	952
Uniprot ID :	P28907
Format :	Purified
Alternative Name :	CD38
Isotype :	Mouse IgG1
Immunogen Information :	Human thymocytes in foetus

Description

CD38 (NAD⁺ glycohydrolase) is a type II transmembrane glycoprotein able to induce activation, proliferation and differentiation of mature lymphocytes and mediate apoptosis of myeloid and lymphoid progenitor cells. Another role of CD38 is provided by enzymatic activity of its extracellular part. CD38 acts as NAD⁺ glycohydrolase converting NAD⁺ into ADP-ribose, as ADP-ribosyl cyclase producing cADPR and as cADPR hydrolase, thus affecting levels of calcium-mobilizing metabolites. ADPR produced by CD38 serves as an important second messenger of neutrophil and dendritic cell migration.

Product Info

Amount :	100 tests
Purification :	Purified by protein-A affinity chromatography
Content :	0.4 ml in PBS Azide
Storage condition :	Store at 2-8°C. Do not freeze.

Application Note

Flow cytometry: *The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10⁶ cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests.*




Figure 1: Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human CD38 (HIT2) PE-Cy™ 5 antibody (4 µl reagent / 100 µl of peripheral whole blood).




Figure 2: Separation of human monocytes (red-filled) from CD38 negative lymphocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human CD38 (HIT2) PE-Cy[™]5 antibody (4 μ l reagent / 100 μ l of peripheral whole blood).