

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

30-2196AF488: Alexa Fluor 488 Conjugated Anti-CD105 / Endoglin Monoclonal Antibody (Clone: MEM-226)

Clonality: Monoclonal
Clone Name: MEM-226
Application: FACS,WB
Reactivity: Rat,Human
Gene: ENG
Gene ID: 2022

Alternative Name: ENG,END,Endoglin,HHT1,ORW1

P17813

Isotype: Mouse IgG2a

Immunogen Information: Recombinant Vaccinia virus containing the human CD105 cDNA.

Description

Uniprot ID:

CD105 (Endoglin) is a homodimeric transmembrane glycoprotein serving in presence of TGFbetaR-2 as a receptor for TGFbeta-1 and TGFbeta-3. CD105 is highly expressed on endothelial cells and promotes angiogenesis during wound healing, infarcts and in a wide range of tumours and its gene expression is stimulated by hypoxia. CD105 prevents apoptosis in hypoxic endothelial cells and also antagonises the inhibitory effects of TGFbeta-1 on vascular endothelial cell growth and migration. Normal cellular levels of CD105 are required for formation of new blood vessels.

Specificity: The antibody MEM-226 reacts with an extracellular epitope of CD105 (Endoglin), a 90 kDa type I homodimerizing membrane glycoprotein expressed on vascular endothelial cells (small and large vessels), activated monocytes and tissue macrophages, stromal cells of certain tissues including bone marrow, pre-B lymphocytes in fetal marrow and erythroid precursors in fetal and adult bone marrow; it is also present on syncytiotrophoblast on placenta throughout pregnancy.

Product Info

Amount: 100 tests

Purified antibody is conjugated with Alexa Fluor 488 NHS ester under optimum conditions and

unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.

Content: Formulation: Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide

Storage condition: Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.

Application Note

Flow cytometry: The reagent is designed for analysis of human blood cells using 4 μ l reagent / 100 μ l of whole blood or 106 cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 1: Flow cytometry analysis (surface staining) of HUVEC (human umbilical vein endothelial cells) and HEK-293 cells with anti-CD105 (MEM-226) Alexa Fluor 488.