

**30-1185PE-Cy5: PE-Cy5 Conjugated Anti-CD34 / Mucosialin Monoclonal Antibody
(Clone:4H11[APG])**

Clonality :	Monoclonal
Clone Name :	4H11[APG]
Application :	FACS
Reactivity :	Human
Conjugate :	R-PE/CY5
Gene :	CD34
Gene ID :	947
Uniprot ID :	P28906
Isotype :	Mouse IgG1
Immunogen Information :	Permanent human cell line derived from peripheral leucocytes of a patient suffering from chronic myeloid leukaemia.

Description

CD34 is a highly glycosylated monomeric 111-115 kDa surface protein, which is present on many stem cell populations. It is a well established stem cell marker, though its expression on human hematopoietic stem cells is reversible. CD34 probably serves as a surface receptor that undergoes receptor-mediated endocytosis and regulates adhesion, differentiation and proliferation of hematopoietic stem cells and other progenitors. CD34 expression is likely to represent a specific state of hematopoietic development that may have altered adhering properties with expanding and differentiating capabilities in both in vitro and in vivo conditions.

Product Info

Amount :	100 tests
Purification :	Purified antibody is conjugated with activated tandem dye of R-phycoerythrin-cyanine 5 (PE-Cy5) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.
Content :	Formulation: Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Storage condition :	Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.

Application Note

Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 106 cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests.




Figure 1: Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human CD34 (4H11[APG]) PE-Cy5 antibody (4 µl reagent / 100 µl of peripheral whole blood).




Figure 2: Separation of human CD34 positive stem cells (red-filled) from lymphocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human CD34 (4H11[APG]) PE-Cy5 antibody (4 µl reagent / 100 µl of peripheral whole blood).