# **w** abeomics

### 30-2925PE-Cy7: Anti-Human CD226 Monoclonal Antibody (Clone:11A8) PE-Cy7 Conjugated

Clonality :	Monoclonal
Clone Name :	11A8
Application :	FACS
Reactivity :	Human, Non-Human Primates
Conjugate :	PE/CY7
Gene ID :	10666
Uniprot ID :	Q15762
Format :	Purified
Alternative Name :	DNAM1, PTA1, TLISA1
Isotype :	Mouse IgG1 kappa

#### Description

CD226 is a type I transmembrane glycoprotein, which is expressed after activation on NK cells, platelets, monocytes, and some T cells. Its expression is increased in patients suffering from some autoimmune diseases or viral infections. CD226 is being phosphorylated on its cytoplasmic tail and interacts with LFA-1 complex (CD11a/CD18), resulting in calcium-independent intercellular contacts. Ligands for CD226 are CD155 and CD112.

#### **Product Info**

Amount :	100 tests
Purification :	Purified by protein-A affinity chromatography
Content :	Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Storage condition :	Store at 2-8°C. Do not freeze.

#### **Application Note**

Flow cytometry: The reagent is designed for analysis of human blood cells using 4  $\mu$ l reagent / 100  $\mu$ l of whole blood or 106 cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests.



Figure 1: Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human CD226 (11A8) PE-Cy<sup>™</sup>7 antibody

For Research Use Only. Not for use in diagnostic/therapeutics procedures.

# **∗** abeomics

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com



Figure 2: Flow cytometry multicolor surface staining pattern of human lymphocytes stained using anti-human CD3 (UCHT1) APC antibody (10  $\mu$ l reagent / 100  $\mu$ l of peripheral whole blood) and anti-human CD226 (11A8) PE-Cy<sup>TT</sup> 7 antibody

Figure 3: Separation of human CD45 negative blood debris (red-filled) from CD226 negative neutrophil granulocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human CD226 (11A8) PE-Cy<sup>™</sup>7 antibody