

### 30-2905PE: PE Anti-Hu CD141 Mab (M80)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	M80
<b>Application :</b>	FACS
<b>Reactivity :</b>	Human
<b>Conjugate :</b>	PE
<b>Gene :</b>	CD141
<b>Gene ID :</b>	7056
<b>Uniprot ID :</b>	P07204
<b>Alternative Name :</b>	Flow cytometry: Recommended dilution: 2-4 µg/ml.
<b>Isotype :</b>	Mouse IgG1
<b>Immunogen Information :</b>	MV4-11 cell line

#### Description

CD141, also known as thrombomodulin or fetomodulin, is a single chain type I transmembrane glycoprotein serving as a receptor for thrombin and as an important cofactor in the protein C anticoagulant system, but it is also involved in embryonic and atherosclerotic plaque development. CD141 is expressed mainly on macrophages, monocytes, a subpopulation of myeloid dendritic cells, on platelets and endothelial cells, but also e.g. on keratinocytes (epithelium). After binding to thrombin, CD141 activates protein C, which degrades clotting factors Va and VIIIa, and as a consequence the amount of thrombin is reduced. Mutations in the CD141 gene can cause a thromboembolic disease known as inherited thrombophilia.

#### Product Info

<b>Amount :</b>	100 Tests
<b>Purification :</b>	Purified antibody is conjugated with R-phycoerythrin (PE) under optimum conditions. Unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.
<b>Content :</b>	Formulation: Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
<b>Storage condition :</b>	Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.

#### Application Note

Flow cytometry: The reagent is designed for analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 106 cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.




Figure 1: Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human CD141 (M80) PE antibody (10  $\mu$ l reagent / 100  $\mu$ l of peripheral whole blood).




Figure 2: Separation of human monocytes (red-filled) from lymphocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human CD141 (M80) PE antibody (10  $\mu$ l reagent / 100  $\mu$ l of peripheral whole blood).