

30-1484PECy7: PE-Cy7 Conjugated Anti-EGFR Monoclonal Antibody (Clone:EGFR1)

| | |
|--------------------------------|--------------------------------------|
| Clonality : | Monoclonal |
| Clone Name : | EGFR1 |
| Application : | FACS |
| Reactivity : | Human |
| Conjugate : | PE/CY7 |
| Gene : | EGFR |
| Gene ID : | 1956 |
| Uniprot ID : | P00533 |
| Format : | Purified |
| Alternative Name : | EGFR,ERBB,ERBB1,HER1 |
| Isotype : | Mouse IgG2b |
| Immunogen Information : | Human epidermoid carcinoma line A431 |

Description

The oncoprotein EGFR (epidermal growth factor receptor), also known as HER1 / ErbB1, is a member of ErbB family of cell surface receptor tyrosine kinases. This 170 kDa transmembrane glycoprotein is often associated with cancerogenesis, although its activation state is controlled at various levels including trafficking and degradation steps. Binding of receptor-specific ligands to the EGFR ectodomain results in formation of homodimeric or heterodimeric kinase-active complexes into which HER2 / ErbB2 is recruited as a preferred partner. Subsequent signaling cascades such as RAS/MAPK and PI3K/AKT pathways lead to cell proliferation and survival responses.

Product Info

| | |
|----------------------------|--|
| Amount : | 100 Tests |
| Purification : | Purified antibody is conjugated with activated tandem dye of R-phycoerythrin-cyanine 7 (PE-Cy™ 7) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography. |
| Storage condition : | Store at 2-8°C protected from light. Do not freeze. |

Application Note

Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10⁶ cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests.

Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10⁶ cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests.




Figure 1: Separation of A431 cells stained using anti-EGFR (EGFR1) PE-Cy™ 7 antibody (concentration in sample 1.67 µg/ml, red-filled) from A431 cells stained using mouse IgG1 isotype control (MOPC-21) PE-Cy™ 7 antibody (concentration in sample 1.67 µg/ml, same as EGFR PE-Cy™ 7 concentration, black-dashed) in flow cytometry analysis (surface staining).