

30-1517AC: APC Conjugated Anti-DLL4 (Clone : MHD4-46)

Clonality :	Monoclonal
Clone Name :	MHD4-46
Application :	FACS
Reactivity :	Human
Conjugate :	APC
Gene :	DLL4
Gene ID :	54567
Uniprot ID :	Q9NR61
Alternative Name :	Delta like ligand 4, AOS6, canonical Notch ligand 4,delta like canonical Notch ligand 4
Isotype :	Mouse IgG1 kappa
Immunogen Information :	recombinant soluble human DLL4

Description

DLL4 (Delta-like 4) is one of five ligands of Notch receptors. It interacts with Notch1 and Notch4. DLL4 is up-regulated at sites of physiologic and pathologic angiogenesis, whereas its expression is low in most adult normal tissues. It is also highly expressed in human clear-cell renal carcinomas, bladder cancers, and breast cancers. Blocking the DLL4-Notch interaction seems to be a promising therapeutic approach.

Specificity : The mouse monoclonal antibody MHD4-46 recognizes the extracellular domain of DLL4 (Delta-like ligand 4), a type I transmembrane protein which plays an important role in vascular development.

Product Info

Amount :	0.1 mg
Purification :	Purified antibody is conjugated with activated allophycocyanin (APC) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.
Content :	Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Storage condition :	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light.

Application Note

Flow cytometry: Recommended dilution: 1-5 µg/ml.




Figure 1 : Separation of DLL4 transfected CHO cells stained using anti-DLL4 (MHD4-46) APC antibody (concentration in sample 5 µg/ml, red-filled) from DLL4 transfected CHO cells stained using mouse IgG1 isotype control (MOPC-21) APC antibody (concentration in sample 5 µg/ml, same as DLL4 APC antibody concentration, black-dashed) in flow cytometry analysis (surface staining).