

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 30-2726PE: PE Conjugated Anti-Hu NKp80 (Clone: 5D12)

Clone Name: 5D12
Application: FACS

**Reactivity:** Monkey, Human

Conjugate: PE
Gene: KLRF1
Gene ID: 51348
Uniprot ID: Q9NZS2

Alternative Name: killer cell lectin like receptor F1 KLRF1, CLEC5C

**Isotype:** Mouse IgG1 kappa

Immunogen Information: recombinant human NKp80 extracellular domain

## **Description**

NKp80, also known as CLEC5C or KLRF1, is a type II transmembrane glycoprotein of the C lectin family, which is expressed in 80 kDa homodimers on NK cells, and subsets of CD8+ alpha/beta T cells, and gamma/delta T cells. It belongs to the activating coreceptors, which induce cytotoxicity, and production of pro-inflammatory cytokines. Its ligand AICL is expressed on myeloid cells.

Specificity: The mouse monoclonal antibody 5D12 recognizes an extracellular epitope of human NKp80 (CLEC5C), a C-type lectin family member, expressed on NK cells and subsets of T cells.

## **Product Info**

Amount: 100 Tests

**Purification :** Purified antibody is conjugated with R-phycoerythrin (PE) under optimum conditions.

Unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.

**Content:** Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide

**Storage condition :** Store at 2-8°C. Do not freeze.

## **Application Note**

Flow cytometry: The reagent is designed for analysis of human blood cells using 10  $\mu$ l reagent / 100  $\mu$ l of whole blood or 10<sup>6</sup> cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.




Figure 1: Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human NKp80 (5D12) PE antibody (10 µl reagent / 100 µl of peripheral whole blood).



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2: Separation of human NKp80 positive lymphocytes (red-filled) from NKp80 negative lymphocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human NKp80 (5D12) PE antibody (10  $\mu$ l reagent / 100  $\mu$ l of peripheral whole blood).