

**12-9281B: Biotinylated Anti-CSF1R antibody(DM107); Rabbit mAb**

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	DM107
<b>Application :</b>	ELISA; Flow Cyt
<b>Reactivity :</b>	Human
<b>Gene :</b>	CSF1R
<b>Uniprot ID :</b>	P07333
<b>Alternative Name :</b>	CSF1R;C-FMS;CD115;CSFR;FIM2;FMS;M-CSFR
<b>Isotype :</b>	IgG

**Description**

The protein encoded by this gene is the receptor for colony stimulating factor 1; a cytokine which controls the production; differentiation; and function of macrophages. This receptor mediates most if not all of the biological effects of this cytokine. Ligand binding activates the receptor kinase through a process of oligomerization and transphosphorylation. The encoded protein is a tyrosine kinase transmembrane receptor and member of the CSF1:PDGF receptor family of tyrosine-protein kinases. Mutations in this gene have been associated with a predisposition to myeloid malignancy. The first intron of this gene contains a transcriptionally inactive ribosomal protein L7 processed pseudogene oriented in the opposite direction. Alternative splicing results in multiple transcript variants. Expression of a splice variant from an LTR promoter has been found in Hodgkin lymphoma (HL); HL cell lines and anaplastic large cell lymphoma.

**Product Info**

<b>Amount :</b>	10 µg / 100 µg
<b>Purification :</b>	Purified from cell culture supernatant by affinity chromatography
<b>Content :</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage condition :</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

**Application Note**

ELISA 1:5000-10000; Flow Cyt 1:100