

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

12-9428B: Biotinylated Anti-UPA antibody(DMC440); IgG1 Chimeric mAb

Clonality: Monoclonal
Clone Name: DMC440
Application: Flow Cyt
Reactivity: Human
Gene: UPA
Uniprot ID: P00749

Alternative Name: ATF; BDPLT5; QPD; u-PA; UPA; URK

Isotype: lgG1

Description

This gene encodes a secreted serine protease that converts plasminogen to plasmin. The encoded preproprotein is proteolytically processed to generate A and B polypeptide chains. These chains associate via a single disulfide bond to form the catalytically inactive high molecular weight urokinase-type plasminogen activator (HMW-uPA). HMW-uPA can be further processed into the catalytically active low molecular weight urokinase-type plasminogen activator (LMW-uPA). This low molecular weight form does not bind to the urokinase-type plasminogen activator receptor. Mutations in this gene may be associated with Quebec platelet disorder and late-onset Alzheimer's disease. Alternative splicing results in multiple transcript variants; at least one of which encodes an isoform that is proteolytically processed.

Product Info

Amount : $10 \mu g / 100 \mu g$

Purification: Purified from cell culture supernatant by affinity chromatography

Content: Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before

lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended

Storage condition: for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient temperature.

Application Note

Flow Cyt 1:100