

36-1791: Monoclonal Antibody to PGP9.5 / UchL1 (pan-Neuronal Marker)(Clone : UCHL1/775)

Clonality :	Monoclonal
Clone Name :	UCHL1/775
Application :	WB,IHC
Reactivity :	Human, Rat
Gene :	UCHL1
Gene ID :	7345
Uniprot ID :	P09936
Format :	Purified
Alternative Name :	UCHL1
Isotype :	Mouse IgG1, kappa
Immunogen Information :	Recombinant human UCHL1 protein

Description


This MAb reacts with a protein of 20-30kDa, identified as PGP9.5, also known as ubiquitin carboxyl-terminal hydrolase-1 (UchL1). Initially, PGP9.5 expression in normal tissues was reported in neurons and neuroendocrine cells but later it was found in distal renal tubular epithelium, spermatogonia, Leydig cells, oocytes, melanocytes, prostatic secretory epithelium, ejaculatory duct cells, epididymis, mammary epithelial cells, Merkel cells, and dermal fibroblasts. Furthermore, immunostaining for PGP9.5 has been shown in a wide variety of mesenchymal neoplasms as well. A mutation in PGP9.5 gene is believed to cause a form of Parkinson's disease.

Product Info


Amount :	100 µg
Purification :	Affinity Chromatography
Content :	100 µg in 500 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic.
Storage condition :	Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid repeated freeze and thaw cycles.

Application Note

Western Blot (1-2ug/ml);Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);



Formalin-fixed, paraffin-embedded Rat Testis stained with Pgp9.5 Monoclonal Antibody (UCHL1/775).



Formalin-fixed, paraffin-embedded Rat Cerebellum stained with Pgp9.5 Monoclonal Antibody (Uchl1/775).