

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

12-9513B: Biotinylated Anti-FLT3 antibody(19A5), IgG1 Chimeric mAb

Clonality: Monoclonal
Clone Name: 19A5
Application: Flow Cyt
Reactivity: Human
Gene: FLT3
Uniprot ID: P36888

Alternative Name: Flt-3;Flk-2;STK-1;CD135;FLK2;FLT-3

Isotype: lgG1

Description

Flt-3(Receptor-type tyrosine-protein kinase FLT3) is also known as FLK-2(Fetal liver kinase-2), STK-1(Stem cell tyrosine kinase 1), CD135. FLT3 is a cytokine receptor which belongs to the receptor tyrosine kinase class III. Tyrosine-protein kinase that acts as cell-surface receptor for the cytokine FLT3LG and regulates differentiation, proliferation and survival of hematopoietic progenitor cells and of dendritic cells. Promotes phosphorylation of SHC1 and AKT1, and activation of the downstream effector MTOR. Promotes activation of RAS signaling and phosphorylation of downstream kinases, including MAPK1/ERK2 and/or MAPK3/ERK1. Mutations that cause constitutive kinase activity promote cell proliferation and resistance to apoptosis via the activation of multiple signaling pathways.

Product Info

Amount : $10 \mu g / 100 \mu g$

Purification: Purified from cell culture supernatant by affinity chromatography

Content: Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before

lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended

Storage condition: for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient temperature.

Application Note

Flow Cyt 1/100