

36-1807: Monoclonal Antibody to von Willebrand Factor / Factor VIII Related-Ag (Endothelial Marker)(Clone : IIIE2.34)

Clonality :	Monoclonal
Clone Name :	III E2.34
Application :	IP,FACS,IFWB,IHC
Reactivity :	Human
Gene :	VWF
Gene ID :	7450
Uniprot ID :	P04275
Format :	Purified
Alternative Name :	VWF,F8VWF
Isotype :	Mouse IgG1, kappa
Immunogen Information :	Recombinant human vWF fragment spanning aa 845-949

Description


von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposi's sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.

Product Info


Amount :	100 µg
Purification :	Affinity Chromatography
Content :	100 µg in 500 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic.
Storage condition :	Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid repeated freeze and thaw cycles.

Application Note

Immunoprecipitation (0.5-1 g/500ug protein lysate); Flow Cytometry (1-2ug/million cells);,Immunofluorescence (1-2ug/ml); Western Blot (1-2ug/ml); ,Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),



Formalin-fixed, paraffin-embedded human Tonsil stained with vWF Monoclonal Antibody (III E2.34)



Formalin-fixed, paraffin-embedded human Tonsil stained with vWF Monoclonal Antibody (III E2.34)