

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-18626: Human 4-1BB(87-118) Protein, hFc Tag

Gene: 4-1BB **Uniprot ID:** 007011

ILA; TNFRSF9; CD137; CDw137; IMD109, Recombinant human 4-1BB(87-118) Protein with C-terminal **Alternative Name:**

Description

The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor contributes to the clonal expansion, survival, and development of T cells. It can also induce proliferation in peripheral monocytes, enhance T cell apoptosis induced by TCR/CD3 triggered activation, and regulate CD28 co-stimulation to promote Th1 cell responses. The expression of this receptor is induced by lymphocyte activation. TRAF adaptor proteins have been shown to bind to this receptor and transduce the signals leading to activation of NF-kappaB. [provided by RefSeg, Jul 2008] Molecular Weight: The protein has a predicted molecular mass of 29.5 kDa after removal of the signal peptide. The apparent molecular mass of 4-1BB(87-118)-hFc is approximately 25-35 kDa due to glycosylation.

Tag: C-Human Fc tag

Product Info

Amount: 50μg / 10μg

The purity of the protein is greater than 90% as determined by SDS-PAGE and Coomassie blue **Purification:**

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before Content:

lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended

Storage condition: for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient temperature.




Figure 1. Human 4-1BB(87-118) Protein, hFc Tag on SDS-PAGE under reducing condition.