

32-1035: ANGPTL4 HEK Recombinant Protein

Alternative Name : ANGPTL4,NL2,ARP4,FIAP,PGAR,HFARP,pp1158,ANGPTL2,Fasting- Induced Adipose Factor,Hepatic Fibrinogen/Angiopoietin-Related Protein,PPARG Angiopoietin-Related Protein.

Description

Source : HEK293. The ANGPTL4 Human Recombinant is manufactured with C-terminal fusion of 11 amino acid FLAG Tag. The ANGPTL4 Flag -Tagged Fusion Protein is a 44.2kDa protein containing 392 amino acid residues of the Angiopoietin-like Protein 4 and 11 additional amino acid residues - Flag Tag . The fasting-induced adipose factor (FIAP, ANGPTL4, PGAR, HFARP) was identified as an adipocytokine up-regulated by fasting, by peroxisome proliferator-activated receptor agonists, and by hypoxia. At the protein level, in human and mouse blood plasma, FIAP was found to be present both as a native protein and in a truncated form. Differentiation of mouse 3T3-L1 adipocytes was associated with the production of truncated FIAP, whereas in human white adipose tissue and SGBS adipocytes, only the native FIAP could be detected. Interestingly, the truncated FIAP was produced by human liver. Experimental data suggest that FIAP is mainly presented in human blood plasma in a truncated form (FIAP-S2), whose level is increased by fenofibrate treatment. Levels of both truncated and native FIAP showed marked inter individual variation but were not associated with body mass index and were not influenced by prolonged semistarvation.

Product Info

Amount : 10 µg
Purification : Greater than 90% as determined by SDS-PAGE.
Content : Filtered and lyophilized from 0.5mg/ml in 20mM Tris buffer and 50mM NaCl pH-7.5.
Storage condition : Store lyophilized Angiopoietin-like Protein 4 Human recombinant at -20°C. Aliquot the product after reconstitution to avoid repeated freezing/thawing cycles. Reconstituted protein can be stored at 4°C for a limited period of time; it does not show any change after two weeks at 4°C.
Amino Acid : GPVQSKSPRF ASWDEMNVLA HGLLQLGQGL REHAERTRSQ LSALERRLSA CGSACQGTEG
STDLPLAPES RVDPEVLHSL QTQLKAQNSR IQQLFHKVAQ QQRHLEKQHL RIQHLQSQFG LLDHKHLDHE
VAKPARRKRL PEMAQPVDPA HNVSRHLRLP RDCQELFQVG ERQSGLFEIQ PQGSPFLVN
CKMTSDGGWT VIQRRHDGSV DFNRPW EAYK AGFGDPHGEF WLGLEKVHSI TGDRNSRLAV
QLRDWDGNAE LLQFSVHLGG EDTAYSLQLT APVAGQLGAT TVPPSGLSVP FSTWDQDHDL
RRDKNCAKSL SGGWWFGTCS HSNLNGQYFR SIPQQRQKLK KGIFWKTWRG RYYPLQATTM
LIQPM AAEAA SAAADYKDDDDK.

Application Note

Add pyrogen free water to a working concentration of 0.5mg/ml and let the lyophilized pellet dissolve completely. Product is not sterile! Please filter the product by an appropriate sterile filter before using it in the cell culture.

