

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-1105: bBTC Recombinant Protein

Description

Source: Escherichia Coli. Betacellulin Bovine Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 80 amino acids and having a molecular mass of 9003 Dalton. Betacellulin Bovine Recombinant is purified by proprietary chromatographic techniques. Btc is a potent mitogen for retinal pigment epithelial cells and vascular smooth muscle cells. The effects of betacellulin are probably mediated by the egf receptor and other related receptors.

Product Info

Amount : 20 μg

Purification : Greater than 95.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

Content: The Betacellulin Bovine Recombinant was lyophilized after extensive dialysis against 50mM

acetic acid.

Lyophilized Betacellulin Bovine Recombinant although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution BTC Bovine should be stored at

4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended

to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Amino Acid: The sequence of the first five N-terminal amino acids was determined and was found to be Asp-

Gly-Asn-Ser-Thr.

Application Note

Storage condition:

It is recommended to reconstitute the lyophilized BTC Bovine in sterile $18M\tilde{A} \square \hat{A} \odot$ -cm H2O not less than $100\tilde{A} \square \hat{A} \mu g/ml$, which can then be further diluted to other aqueous solutions. The ED50, calculated by the dose-dependent proliferation of murine BALBC 3T3 cells (measured by 3H-thymidine uptake) is < 10.0 ng/ml, corresponding to a Specific Activity 100,000 units/mg.

