

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-1107: mCDNF Recombinant Protein

Alternative Name : Cerebral dopamine neurotrophic factor,ARMET-like protein 1,Conserved dopamine neurotrophic factor,Cdnf,Armetl1,9330140G23.

Description

Source: Escherichia Coli. CDNF Mouse Recombinant produced in E.coli is a single, non-glycosylated polypeptide chain containing 163 amino acids and having a molecular mass of 18.5kDa.The CDNF is purified by proprietary chromatographic techniques. CDNF is a member of the ARMET family and acts as a trophic factor for dopamine neurons. CDNF inhibits the 6-hydroxydopamine (6-OHDA)-induced degeneration of dopaminergic neurons. When CDNF controlled after 6-OHDA-lesioning, it reestablishes the dopaminergic function and inhibits the degeneration of dopaminergic neurons in substantia nigra. CDNF is universally expressed in neuronal and non-neuronal tissues. The highest levels in the brain are found in the optic nerve and corpus callosum.

Product Info

Amount: 20 µg

Purification: Greater than 97.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

Content: CDNF protein was lyophilized from a 0.2µm filtered concentrated solution in 1xPBS, pH 7.4.

Lyophilized CDNF although stable at room temperature for 3 weeks, should be stored

Storage condition:

desiccated below -18°C. Upon reconstitution CDNF should be stored at 4°C between 2-7 days

and for future use below -18 $^{\circ}$ C. For long term storage it is recommended to add a carrier

protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Amino Acid : QGLEAGVGPR ADCEVCKEFL DRFYNSLLSR GIDFSADTIE KELLNFCSDA KGKENRLCYY LGATTDAATK

ILGEVTRPMS VHIPAVKICE KLKKMDSQIC ELKYGKKLDL ASVDLWKMRV AELKQILQRW GEECRACAEK

SDYVNLIREL APKYVEIYPQ TEL.

Application Note

It is recommended to reconstitute the lyophilized CDNF in sterile 18M-cm H2O not less than $100\tilde{A} \square \hat{A} \mu g/ml$, which can then be further diluted to other aqueous solutions. CDNF Mouse is able to enhance neurite outgrowth of E16-E18 rat embryonic cortical neurons when immobilized at 5-30 $\tilde{A} \square \hat{A} \mu g/mL$ on a nitrocellulose-coated microplate.

